
Chairside Preparation Guide

ANTERIOR CROWN PREPARATION 1.0-1.5 mm reduction at gingival margin 1.5 mm lingual contact clearance reduction 1.5 mm to 2.0 mm incisal reduction Football shaped blur used for lingual reduction.



VENEER PREPARATION GUIDE

FACIAL REDUCTION

A medium grit, round-ended diamond bur is used to remove a uniform thickness of facial enamel by joining the depth-cut grooves.




• Uniform facial preparation is recommended.

CHAMFER MARGIN

Correct preparation of the chamfer margins interproximally allows the appropriate bulk of porcelain.







1.0 – 1.5 mm

1.0 – 1.5 mm

1.0 mm


ecommended preparation for anterior teeth

Recommended preparation for posterior teeth

- The Ideal Preparartion should include a Shoulder or Chamfer to allow for a continous and clearly visible precise detectable margin.
- Bevel margins should be avoided.
- All preparations should have a smooth rounded surface with no sharp angles.
- A double retraction cord technique for impressions is highly reccommended.

INCISAL REDUCTION

The diamond bur is angled to bevel back the incisal edge.



CONTACT REDUCTION

There are two choices for interproximal margins:

- 1. Interproximal margins should remain facial to the natural enamal contact.
- 2. Prepare through the natural contact and extend the margin to the lingual side.
- When closing a diastema interproximal margins need to extend to the lingual marginal ridges, eliminating any and all undercuts.

Have a question? Call and ask for Bob Marbach, BA, CDT.

800-683-1025 210-735-1433

